

PATIENT-SIGNED HIPAA/DATA PRIVACY CONSENT:

By signing this authorization, I authorize my health plans, physicians and staff, other healthcare providers, and pharmacy providers (collectively, my "Providers") to disclose information, including but not limited to, personal health information about me or my minor child, including information related to my or my child's medical condition, treatment, care management, and health insurance coverage and claims, any prescription (including fill/refill information), as well as information provided on this form (collectively, "Personal Health Information"), to CSL Behring and its representatives, agents, and contractors, including CSL Behring's support program(s) (collectively "CSL Behring Entities") for the purposes of:

- (1) establishing eligibility for insurance benefits including but not limited to coverage for prescription drugs;
- (2) evaluation and enrollment in one or more financial assistance program(s) offered by CSL Behring Entities, such as a co-pay mitigation program and/or patient assistance programs (if one or more of such programs apply to my treatment with a CSL Behring therapy);
- (3) enrollment in available patient services programs offered by CSL Behring Entities;
- (4) communication about my treatment with me or my Providers, including by contacting me directly to facilitate the dispensing of medication and scheduling shipments and refill reminders;
- (5) providing product support and adherence services through CSL Behring Entities;
- (6) evaluating the effectiveness of CSL Behring's support program(s); and
- (7) any other related support, education, and assistance services related to my treatment with CSL Behring therapy and/or living with my disease (collectively, the "Services").

Further, I authorize any of the CSL Behring Entities to contact me by mail, telephone, or e-mail for relevant follow-up to any of the aforementioned services. CSL Behring entities include but are not limited to brand specific support through hub service providers, pharmacy service providers, nurse self-infusion training providers and/or nurse adherence providers. I understand that these CSL Behring Entities may collect Personal Health Information from me for the purposes listed above, and that such collection is subject to CSL Behring's Privacy Policy.

I understand that my pharmacy Providers, including those Providers who dispense free trials or commercially-reimbursed doses of CSL Behring products, may disclose to the CSL Behring Entities certain Personal Health Information regarding the dispensing of my prescription and that such disclosure will result in remuneration to my pharmacy Provider(s). If necessary or if requested by my prescriber, I authorize CSL Behring Entities to forward my prescription to a dispensing pharmacy on my behalf.

I understand that once my Personal Health Information is disclosed to the CSL Behring Entities under this authorization, it may no longer be protected by state and/or federal privacy laws and may be further disclosed by the CSL Behring Entities. However, I understand that the CSL Behring Entities will disclose my Personal Health Information only for the limited purposes described above, or as I may further authorize in writing, or as permitted or required by law.

MARKETING OPT IN Receive relevant CSL Behring communications and/or patient resources (Optional)

I further authorize the CSL Behring Entities to provide my information and/or contact me regarding education, training, and ongoing support on the use of CSL Behring therapy and that may be of interest to me. I understand that the CSL Behring Entities may contact me by mail, telephone/text message, or email.

I understand that, in some cases, CSL Behring uses third-parties under contract with CSL Behring to administer its educational and patient benefit programs. I understand that when contacting me on behalf of CSL Behring about these programs, these third-parties will have access to my email address, address, and/or phone number. I understand that, like CSL Behring, these third-parties will not rent, sell, publicly post, or share my personal information.

Important Safety Information

Hizentra is indicated for:

- Treatment of primary immunodeficiency (PI) in adults and pediatric patients 2 years and older.
- Maintenance therapy in adults with chronic inflammatory demyelinating polyneuropathy (CIDP) to prevent relapse of neuromuscular disability and impairment.
 - Limitation of use: maintenance therapy in CIDP has been systematically studied for 6 months and for a further 12 months in a follow-up study. Continued maintenance beyond these periods should be individualized based on patient response and need for continued therapy.

For subcutaneous infusion only.

WARNING: Thrombosis may occur with immune globulin products, including Hizentra. Risk factors may include: advanced age, prolonged immobilization, hypercoagulable conditions, history of venous or arterial thrombosis, use of estrogens, indwelling vascular catheters, hyperviscosity, and cardiovascular risk factors.

For patients at risk of thrombosis, administer Hizentra at the minimum dose and infusion rate practicable. Ensure adequate hydration in patients before administration. Monitor for signs and symptoms of thrombosis and assess blood viscosity in patients at risk for hyperviscosity.

Hizentra is contraindicated in patients with a history of anaphylactic or severe systemic reaction to human immune globulin (Ig) or components of Hizentra (eg, polysorbate 80), as well as in patients with immunoglobulin A deficiency with antibodies

I understand that data related to my enrollment in any CSL Behring program may be collected, analyzed and shared among CSL Behring Entities. I also understand that CSL Behring Entities, which operate the patient support services for CSL Behring, may receive compensation from CSL Behring in connection with the patient support services.

I understand that I may refuse to sign this authorization. I understand, however, that if I do not sign this authorization, I may not be able to receive patient support services through CSL Behring. I understand that my treatment with a CSL Behring therapy (other than participation in a free trial program), payment for treatment, insurance enrollment, or eligibility for insurance benefits are not conditioned upon my agreement to sign this authorization. I understand that patient support services provided by CSL Behring are not insurance and that CSL Behring has the right to rescind, revoke or amend any service at any time without notice.

I understand that I am entitled to a copy of this authorization.

I understand that if CSL loans me durable medical equipment or other medical equipment through the Services, CSL reserves the right to seek reimbursement from me for all unreturned DME or equipment.

I understand that I may change my mind and cancel this authorization at any time by writing a letter requesting such cancellation to CSL Behring c/o Patient Services P.O. Box 61501 King of Prussia, PA 19406 or by calling the CSL Behring Customer Affairs toll free number 1-888-508-6978 and that this cancellation will end my participation in CSL Behring Services and will not apply to any information already used or disclosed through this authorization before notice of the cancellation is received by CSL Behring Entities. This authorization expires five (5) years from the date signed below, or earlier, if required by state law. CSL Behring will not retain this data beyond the maximum period allowed by law.

Under certain circumstances, by law you may have certain rights regarding CSL Behring's use of your data. You may have the right to receive information about what data CSL Behring has collected about you. You may have the right to ask CSL Behring to delete certain personal information about you, but only when CSL Behring does not have a legal reason for retaining your personal information. If you exercise these rights, you will be asked to verify your identity, if someone else will exercise your rights on your behalf, they will need to prove that they have your permission to do so. To exercise your rights, please contact us through <https://privacyinfo.csl.com/> or toll free by phone at (833) 704-0018. For more information about how CSL Behring handles personal information, please visit our privacy policy at <https://www.cslplasma.com/privacy-policy>.

PATIENT OR PARENT/GUARDIAN AUTHORIZATION SIGNATURE:

Date _____

RELATIONSHIP TO PATIENT (IF APPLICABLE): _____

If I change my mind in the future and do not wish to receive information related to a CSL Behring therapy, or any related products or services, or to be contacted occasionally for market research purposes, I understand that I may change my mind and cancel this authorization at any time by writing a letter requesting such cancellation to CSL Behring c/o Patient Services P.O. Box 61501 King of Prussia, PA 19406 or by calling the CSL Behring Customer Affairs toll free number 1-888-508-6978. This cancellation will not apply to any information already used or disclosed through this authorization.

PATIENT OR PARENT/GUARDIAN AUTHORIZATION SIGNATURE:

Date _____

RELATIONSHIP TO PATIENT (IF APPLICABLE): _____

against IgA and a history of hypersensitivity. Because Hizentra contains L-proline as stabilizer, use in patients with hyperprolinemia is contraindicated.

IgA-deficient patients with anti-IgA antibodies are at greater risk of severe hypersensitivity and anaphylactic reactions. Thrombosis may occur following treatment with Ig products, including Hizentra.

Monitor patients for aseptic meningitis syndrome (AMS), which may occur following treatment with Ig products, including Hizentra. In patients at risk of acute renal failure, monitor renal function, including blood urea nitrogen, serum creatinine and urine output. In addition, monitor patients for clinical signs of hemolysis or pulmonary adverse reactions (eg, transfusion-related acute lung injury [TRALI]).

Hizentra is derived from human blood. The risk of transmission of infectious agents, including viruses and, theoretically, the Creutzfeldt-Jakob disease (CJD) agent and its variant (vCJD), cannot be completely eliminated.

The most common adverse reactions (observed in $\geq 5\%$ of study subjects) were local infusion-site reactions, as well as headache, diarrhea, fatigue, back pain, nausea, extremity pain, cough, upper respiratory tract infection, rash, pruritus, vomiting, upper abdominal pain, migraine, arthralgia, pain, fall, and nasopharyngitis.

The passive transfer of antibodies can interfere with response to live virus vaccines and lead to misinterpretation of serologic test results.

Please see accompanying full prescribing information for Hizentra.

To report SUSPECTED ADVERSE REACTIONS, contact the CSL Behring Pharmacovigilance Department at 1-866-915-6958 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.