

Ship Meds to: ☐ Patient's Home ☐ Prescriber's Office

Prescriber Information

Prescriber Name:		☐ MD ☐ DO ☐ NP ☐ PA		NPI:
Office Contact:		Practice Name / Supervising MD:		
Address:		City:		
State:	Zip:	Phone:	Fax:	

Patient Information | PLEASE SEND COPY OF INSURANCE CARD

Patient's Name:	Last 4 Digits of SS#:	DOB: / /	Sex: ☐ M ☐ F	Weight:	Height:	Diabetic: ☐ Y ☐ N
Address:	City:	State:	Zip:	Allergies:		
Home Phone:	Work Or Cell:	HIPAA Contact:	Emergency #:	Interpreter Needed? ☐ Y ☐ N		

Insurance Information

Primary Insurance:	Policy ID:	Group #:
Policyholder Name:	Policyholder DOB:	BIN: PCN:

Clinical Information | PLEASE SEND COPY OF MEDICAL AND PRESCRIPTION INSURANCE CARDS, PROGRESS NOTES AND LAB REPORTS, SUPPORTING DIAGNOSIS, CO-MORBIDITIES AND LAB VALUES

ICD-10/Diagnosis Code: ☐ Multiple Sclerosis (G35) ☐ Other:	Has patient been previously treated for this condition? ☐ Y ☐ N	
Type: ☐ Clinically isolated syndrome ☐ Relapsing-Remitting ☐ Primary Progressive ☐ Secondary Progressive		
Prior failed medication (medication and duration of treatment/reason for d/c): ☐		
Patient currently on therapy? ☐ Y ☐ N Medication(s):	Will patient be stopping above medication before starting new therapy? ☐ Y ☐ N	Discontinuation Date: / /
Is prescriber a Neurologist? If no, please include neurology consult if available ☐ Y ☐ N	☐ Other:	Number of relapses in past year: Last MRI date: / / Any MRI changes? ☐ Y ☐ N
Is patient pregnant, nursing or planning pregnancy? ☐ Y ☐ N ☐ N/A	☐ Serum Creatinine:	☐ Creatinine Clearance:

Prescription Information

Medication	Dose/Strength	Sig	Quantity	Refills
☐ PLEGRIDY™	Starter Pack: ☐ Prefilled syringe (1x63mcg/0.5ml, 1x94mcg/0.5ml) ☐ Autoinjector pen (1x63mcg/0.5ml, 1x94mcg/0.5ml) ☐ 125mcg/0.5ml PFS ☐ 125mcg/0.5ml autoinjector	Dose Titration: ☐ Inject 63mcg SQ on day 1 and 94mcg SQ on day 14 ☐ Maintenance Dose: Inject 125mcg SQ every 14 days, starting on day 29	☐ Titration Dose: 28 day supply ☐ Maintenance Dose: 28 day supply	No Refills
☐ REBIF® ☐ REBIF® REBIDOSE® Autoinjector	☐ Titration Pack (8.8mcg/22mcg) (#12) ☐ 22mcg/0.5ml PFS (#12) ☐ 44mcg/0.5ml PFS (#12)	Dose Titration: ☐ Inject 8.8mcg SQ 3x a week at weeks 1-2, 22mcg SQ 3x a week at weeks 3-4, and 44mcg SQ 3x a week at weeks 5+ (48 hours apart) ☐ Inject 4.4mcg SQ 3x a week at weeks 1-2, 11mcg SQ 3x a week at weeks 3-4, and 22mcg SQ 3x a week at weeks 5+ (48 hours apart) ☐ Maintenance: Inject 22mcg (0.5ml) SQ 3x a week (48 hours apart) ☐ Maintenance: Inject 44mcg (0.5ml) SQ 3x a week (48 hours apart) ☐ Other Regimen:	☐ Titration Dose: 28 Day Supply (12 pens or syringes) ☐ Maintenance Dose: 28 Day Supply	
☐ Other Specialty:				

Injection Training

☐ Patient received injection training	☐ Prescriber's office to provide injection training	☐ Meijer to coordinate injection training
---	---	---

By signing this form and utilizing our services, you are authorizing Meijer and its employees to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies.

Prescriber Signature:	Date	Prescriber Signature:	Date
-----------------------	------	-----------------------	------

Substitution Permitted

Dispense as Written

If brand is required, please write "DAW" in the box to the right.